

Diagnostic Test Studies

Understanding Results

Learning objectives:

1. Importance of pre-test probability
2. Sensitivity/Specificity
3. Likelihood Ratio (LR): **MEMORIZE THE DEFINITION**
- it will be on the IRAT
4. The LR is pre-test-probability independent
5. Calculating the post-test probability

1

What a diagnostic test does

Pre-test probability <i>(Probability that the patient has disease prior to administering the test)</i>	x	"Results of diagnostic test"	=	Post-test probability <i>(Probability that the patient has disease given the additional information of the test results)</i>
--	----------	-------------------------------------	----------	--

What is a pre-test probability and where can we find it?

2

Pre-test probability

- Best: prevalence among my patients
- If don't know, then..
 - prevalence noted in the clinical study
 - ask a local expert
 - make an educated guess

A pre-test probability **MUST** be assigned
in order to figure out the post-test probability

3

"Results of diagnostic test"

Sensitivity and Specificity

- Sensitivity is the proportion of people with a disease who test positive
- Specificity is the proportion of people without a disease who test negative

4

Example of a 2x2 table

	Truly sick	Truly well	
Positive test	TP 95	FP 100	195
Negative test	FN 5	TN 800	805
	100	900	1000

5

Sensitivity: How good is the test when you're sick?

	Truly sick	
Positive test	TP 95	<p>This test correctly picks up 95/100 people who are sick.</p> <p>The sensitivity is 95%</p>
Negative test	FN 5	
	100	

6

Specificity: How good is the test when you're healthy?

This test correctly classifies 800/900 people who are healthy.

Its specificity is 89%

Positive test

Negative test

Truly well

	FP
Positive test	100
Negative test	800
	900

7

Using sensitivity/specificity

- Sensitivity and specificity are test characteristics that are **independent** of disease prevalence (pre-test probability)
- With sensitivity, specificity, and *your patient's* pre-test probability, you can compute *your patient's* post-test probability of having the disease
- **One nice way to compute the post-test probability of disease with sensitivity and specificity is with the Likelihood Ratio**

8

Are also Likelihood Ratios independent of disease prevalence (pre-test probability)?

YES!

(LR's are combinations of sensitivity and specificity)

9

Likelihood Ratio (LR)

MEMORIZE THIS AND THINK ABOUT IT!

Definition of LR:

[for any given test result]

“The probability that the patient comes from the sick rather than the healthy population”

10

Likelihood Ratio

MEMORIZE THIS AND THINK ABOUT IT.

For any given test result, "The probability that the patient comes from the sick rather than the well population"

- Each test result (e.g., positive, negative) has a likelihood ratio (LR+, LR-)
 - +LR should be greater than 1
 - -LR should be less than 1 (fractional)
- LR of 1 means the test result adds no new information (result is equally likely to occur in a sick as in a well person)

+LR means the LR for a positive test
-LR means the LR for a negative test

11

What a diagnostic test does

Pre-test "probability" <i>(Probability that the patient has disease prior to administering the test)</i>	X	Likelihood Ratio (Inherent Test Property)	=	Post-test "probability" <i>(Probability that the patient has disease given the additional information of the test results)</i>
--	----------	---	----------	--

12

Calculation of LR's

Notice that the LR is a **combination** of **SENSITIVITY AND SPECIFICITY**

	Truly sick	Truly well	
Positive test	TP 95	FP 100	195
Negative test	FN 5	TN 800	805
	100	900	1000

+LR = $[95/100]/[100/900]$
+LR = $\text{sensitivity}/(1-\text{specificity})$
+ LR = 8.55

-LR = $[5/100]/[800/900]$
-LR = $(1-\text{sensitivity})/\text{specificity}$
-LR = 0.056

+LR means the LR for a positive test
-LR means the LR for a negative test

13

The remainder of the slides discuss:

Calculating the Post-Test Probability from the Pre-Test Probability and LR

the mathematical way

the nomogram way

the online, Dr. Alan Schwartz, way

14


the mathematical way

Pre-test probability (really the odds) <i>(Probability that the patient has disease prior to administering the test)</i>	X	Likelihood Ratio (Inherent Test Property, Prevalence Independent)	=	Post-test probability <i>(Probability that the patient has disease given the additional information of the test results)</i>
---	----------	--	----------	---

- Convert the pre-test probability (prevalence) to the pre-test odds
pre-test odds (Pr) = prevalence / (1 - prevalence)
- Then calculate the post-test odds: **Pr x LR = post-test odds of disease**
- Finally, convert the post-test odds back to a probability
Probability of disease = [post-test odds] / [1 + post-test odds]

15

the nomogram way




Pre Test Probability

LR

Post-Test Probability

16


Diagnostic Test Nomogram



For example, if the pre-test probability is 10% and the +LR is 9 and -LR is 0.05, then the post-test probability for a positive test is ~45% and for a negative test is ~1%

17

Here is spare nomogram



18

the online, Dr. Alan Schwartz, way

<http://ulan.mede.uic.edu/~alanzs/tools.html>

OR Google EBM ALAN – First Hit

(The website will do *all* your calculations)

19

EBM and Decision Tools by Alan Schwartz

Below you will find links to decision-making tools and exercises developed by [Alan Schwartz](#) and used for evidence-based medicine or medical decision making. Each link opens in its own window.

Tools

[Click here](#)

- [Diagnostic Test Calculator](#) - Given a 2x2 table (or prevalence/sens/spec or prevalence/LRs), compute everything else, including confidence intervals and optionally the impact of the test on action thresholds, and display a graphical nomogram. The [Perl source code](#) for the calculator is available under an open source software license. A [mobile version](#) is now available for use on iPhones and other small-screen browsers.
- [NNI/NNII Calculator](#) - Given information about probability of an event under control and experimental treatment, calculate risk increase/decrease and needed to treat or harm, including confidence intervals.

Exercises

- [Diagnostic Test Cutoffs](#) - A graphical demonstration of the effect of changing cutoff scores on sensitivity and specificity of a test.
- [Statistical Testing Thresholds](#) - A graphical demonstration like the above, but written in terms of statistical test theory (type I and II error)
- [Diagnostic test exercise](#) - Test your knowledge about properties of diagnostic tests.
- [Utility Assessment](#) - Assess your utility for an health state using standard gamble, time tradeoff, and rating scale techniques.
- [Multi-attribute Utility Assessment](#) - Assess the utility of pain killers using multiple attributes, weighted by importance. Demonstrates the SMARTER system MAUT.
- [Markov model simulation](#) - Simulates a simple hypothetical markov model for diabetes.
- [Cost-effectiveness perspectives exercise](#) - Perform some analyses of the cost-effectiveness of different breast cancer screening and treatment policies.



This Resource Successfully Peer Reviewed by [MedEdPORTAL](#) on 4/13/06.
MedEdPORTAL Publication Number: 209

Alterations to this Resource Created After This Date Have Not Been Reviewed By MedEdPORTAL

Subsequent Revision	Date or Frequency of Revision
Added link to mobile version of diagnostic test calculator (same mathematical engine, different user interface)	17 February 2012
Added personal action thresholds to diagnostic test calculator	12 November 2007
Added tree diagram of 2x2 table to diagnostic test calculator, suggested by J. Peter Donnelly	21 April 2010

Diagnostic Test Calculator

Fill out one of the sections below on the left, and then click on the 'Compute' button. Sections you don't fill out will be computed for you, and the nomogram on the right will display the probability that a patient has the disease after a positive or negative test.

Numbers of patients with and without the disease who test positive and negative:

	Disease present	Disease absent	Total
Test positive			
Test negative			
Total			

or
 disease prevalence, test sensitivity, and test specificity (and, optionally, sample size):

Prevalence (e.g. 0.10):	
Sensitivity (e.g. 0.80):	
Specificity (e.g. 0.80):	
Total sample size:	

21

Firefox File Edit View Go Bookmarks Tools Window Help Wed 6:28 PM

Diagnostic Test Calculator

This calculator can determine diagnostic test characteristics (sensitivity, specificity, likelihood ratios) and/or determine the post-test probability of disease given given the pre-test probability and test characteristics. Given sample sizes, confidence intervals are also computed.

Fill out one of the sections below on the left, and then click on the 'Compute' button. Sections you don't fill out will be computed for you, and the nomogram on the right will display the probability that a patient has the disease after a positive or negative test.

Numbers of patients with and without the disease who test positive and negative:

	Disease present	Disease absent	Total
Test positive	95	100	
Test negative	5	800	
Total			

or
 disease prevalence, test sensitivity, and test specificity (and, optionally, sample size):

Prevalence (e.g. 0.10):	
Sensitivity (e.g. 0.80):	
Specificity (e.g. 0.80):	
Total sample size:	

or
 disease prevalence, positive likelihood ratio, and negative likelihood ratio (and, optionally, sample size):

Prevalence (e.g. 0.10):	
PLR (e.g. 4):	

22

Firefox File Edit View Go Bookmarks Tools Window Help 40 (Wed 6:21 PM)

Diagnostic Test Calculator

on the right will display the probability that a patient has the disease after a positive or negative test.

Numbers of patients with and without the disease who test positive and negative:

	Disease present	Disease absent	Total
Test positive	95	100	195
Test negative	5	800	805
Total	100	900	1000

Compute

or disease prevalence, test sensitivity, and test specificity (and, optionally, sample size):

Prevalence (e.g. 0.10):	0.100
Sensitivity (e.g. 0.80):	0.95
Specificity (e.g. 0.80):	0.89
Total sample size:	1000

Compute

or disease prevalence, positive likelihood ratio, and negative likelihood ratio (and, optionally, sample size):

Prevalence (e.g. 0.10):	0.100
+LR (e.g. 4):	8.55
-LR (e.g. 0.01):	0.06
Total sample size:	1000

Compute

Clear Entries

Prior: 0.1 (Disease present), 0.9 (Disease absent)
 Likelihood ratio: 19.0 (Disease present / Disease absent)
 Posterior probability: 0.49 (Disease present), 0.51 (Disease absent)

POSITIVE TEST:
 Positive Likelihood ratio: 8.55
 95% confidence interval: [7.07, 10]
 Posterior probability (odds): 49% (1.0)
 95% confidence interval: [44%, 53%]

NEGATIVE TEST:
 Negative Likelihood ratio: 0.06
 95% confidence interval: [0.02, 0.13]
 Posterior probability (odds): 5% (0.0)
 95% confidence interval: [0%, 15%]

Odds = Probability / (1 - Probability)
 +LR = Sensitivity / (1 - Specificity)
 -LR = (1 - Sensitivity) / Specificity
 Posterior Odds = Prior Odds x LR

23